



# Anencephaly

## Equipment

|   |  |    |  |
|---|--|----|--|
| 1 |  | 8  |  |
| 2 |  | 9  |  |
| 3 |  | 10 |  |
| 4 |  | 11 |  |
| 5 |  | 12 |  |
| 6 |  | 13 |  |
| 7 |  | 14 |  |

## Procedure

|   |  |
|---|--|
| 0 | Introduce yourself to the patient Ask the patient's name.  |
| 1 | Describe the findings such as the USG did not find the upper head. Asking for understanding about Anencephaly, taking a history of genetic disease Congenital disease, drug use, anticonvulsant rental |
| 2 | Anencephaly is a spontaneous disease for many reasons, and the fetus is unable to survive after birth. Effects on pregnancy, such as difficult delivery, polyhydramnios                                |
| 3 | There is a 4% chance of recurrence in the previous pregnancy. Folic a 4 mg/day is recommended from 12 weeks before gestation to 12 weeks after gestation.  |
| 4 | Ask about the willingness to terminate the pregnancy.  |
| 5 | Advise on termination of pregnancy, Cytotec or misoprotal vaginal suppositories.   |
| 6 | Complications such as fever, diarrhea, uterine perforation, hemorrhage, encourage the patient to become pregnant.  |
| 7 | Recommended if next time pregnant Should receive pregnancy in the first place, receive folic a, repeat USG, see for fetal abnormalities.   |
| 8 | Don't be rude, explain clearly and politely. Does not speak technical vocabulary / English   |
| 9 | Show interest, make eye contact, nod in response. Give the patient an opportunity to ask questions.  |

