

## Diagnosis and Treatment of Ventricular Tachycardia

Equipment		
1	8	
2	9	
3	10	
4	11	
5	12	
6	13	
7	14	
Procedure		
0	The teacher reported having Ventricular Tachycardia.	
1	The test taker adjusts or tells him to adjust the power of the Defibrillator 100 J (if using higher or lower power is considered incomplete).	
2	Adjust Synchonize mode on, raise the paddle, apply gel, press the button on both sides of the paddle at the same time.	
3	The paddle is placed on the patient's chest at the apex position and at the sternum (right upper parasternal area), with the sternum peddle placed on the right side of the sternum, not on the sternum.	
4	Tell people around you to clear / step away from the robot. Press the buttons on both peddles at the same time. Make a defibrillator to deduct points.	
5	Test taker reported ventricular tachycardia.	
6	The examinee is palpated with a carotid pulse in no more than 10 seconds. Let the assistant tell you that you can feel the pulse)	
7	The test taker or asks the BP tester to adjust or tell him to adjust the power of the machine. Defibrillator increase to 150–200 J	
8	The test taker repeats steps 2-8 completely.	
9	The test taker responds or shows a pulse, informs the blood pressure measurement.	

TEL: 090-979-1212 Website: <u>www.Jia1669.com</u> Line: @jia1669 Fb.com/Jia1669