Non hemorrhagic shock

Initial assessment
- ABCs
- Airway management and C-spine protection
- Assess: LOC, V/S, traumatic shock
- Incident history; Mechanism of injury

Notify Trauma physical/General physician

Cardiogenic/Obstructive Shock
- Dyspnea, trachycardia
- Decreases breath sound
- Trachea deviation
- Unilateral chest movement
- Paradoxical movement

- O_2 mask with reservoir bag 10-12 LPM or other O2 supplement
- **<u>If tension pneumothortax</u>**: prepare for decompression
- Prepare for: definite Airway
- Prepare for: cardiocentesis, thoracocentesis, ICD, surgical treatments
- Fluid management, keep warm
- Prepare for: X-ray, CT, diagnostic tests
- Maintain oxygenation
- Monitor: V/S, N/S, LOC, O_2sat.,EKG, capillary refill time
- Record

Distributive/Spinal Shock
- Paresis
- Paralysis

- O_2 mask with reservoir bag 10-12 LPM
- Fluid management
- Keep warm
- Medication administration
- Prepare for: definite airway
- Prepare for: X-ray, CT, diagnostic tests
- Maintain normotension
- Monitor: V/S, N/S, LOC, O_2sat., capillaryrefill time
- Record

Reassessment

Definite care to OR/ICU/Unit/transfer