

Agaroses

Smart Agarose

Smart Agarose is an agarose ideal for routine rapid separation of DNA and RNA fragments as well as PCR products, the preparation of plasmids, and for screening, cloning and blotting techniques.

Features:

- · Easy dissolution and rapid gelling.
- Excellent transparency and low background staining gives clear band visibility.
- · Sharp and well defined bands.
- Very low DNA binding.

Applications:

- BioMax has high gel strength even at low concentrations, so use rates are 0.75 - 2%.
- It is effective in blotting and in separations of nucleic acid fractions from 250 bp to 23 Kb.

Markers: Lane 1:1Kb ladder Lane 2: 250 bp ladder Lane 3: 100 bp ladder

Clarity 1.5% (NTU)

Gel Strength 1% (g/cm²)

DNase/RNase activity

Order Information:

*FFO (electroendosmosis

Moisture

Ash

EEO*

Sulfate

Clarity 1.5% (NTU)

Gel Strength 1% (g/cm²)

DNase/RNase activity

Gel background

DNA resolution ≥ 1000 bp

Gel Strength 1.5% (g/cm²)

Gelling Temperature 1.5% (°C)

Melting Temperature 1.5% (°C)

Gel Strength 1.5% (g/cm²)

Gelling Temperature 1.5% (°C)

Melting Temperature 1.5% (°C)

Sulfate

Description	Cat No.
Smart Agarose 100 G	SB101-100G
Smart Agarose 500 G	SB101-500G

Specifications & Functional Tests:

Specifications & Functional Tests:

≤ 0.45%

≤ 0.15%

≥ 1000

≥ 2000 36 ± 1.5

 88 ± 1.5

D-1LE/QS

≤ 10%

≤ 0.4%

≤ 0.1%

≥ 1200

≥ 2500

 36 ± 1.5

 88 ± 1.5

Very low

None detected

Finely resolved

≤3

0.05-0.13

None detected

≤4

LE Agarose

Low EEO Agarose, High electrophoresis mobility

Suitable for:

- Nucleic acid analytical of fractions from 250 bp to 23 Kb
- Preparative electrophoresis
- Blotting
- Protein electrophoresis ex. radial immunodiffusion

- Extraordinary mechanical resistance for more reliable and easier handling.
- Possibility of varying pore size in accordance with particle size by modifying the gel concentration.
- Easy preparation of the gel by simple dilution in aqueous buffers either by standard boiling or microwaving.
- · Greater thermal stability due to high hysteresis (difference between gelling and melting temperatures).
- Excellent transparency of the gel and high visibility.
- · Exceptionally low absorption of staining agents.
- Absence of toxicity (polyacrylamide is neurotox

ES. DXIC).			Order Information:			
JXIC	_			Description	Cat No.	
A B			LE Agarose 100 G	SB102-100G		
				LE Agarose 500 G	SB102-500G	
	11 11	- -				
-			D1-I	-LE Agarose gels in 1X TAE buffer		













A-0.75%, B-1%, C-1.25% Marker: 1Kb ladder. Electrophoresis conditions: submarine gel, 2 hours 30 min, 4.5 V/cm in 1X TAE buffer.